arXiv:math/0602099v1 [math.DS] 6 Feb 2006

A COUNTEREXAMPLE TO A MULTIDIMENSIONAL VERSION OF THE WEAKENED HILBERT'S 16-TH PROBLEM

MARCIN BOBIEŃSKI AND HENRYK ŻOŁĄDEK

Abstract. In the weakened 16th Hilbert's Problem one asks for a bound of the number of limit cycles which appear after a polynomial perturbation of a planar polynomial Hamiltonian vector field. It is known that this number is finite for an individual vector field. In the multidimensional generalization of this problem one considers polynomial perturbation of a polynomial vector field with invariant plane supporting a Hamiltonian dynamics. We present an explicit example of such perturbation with infinite number of limit cycles which accumulate at some separatrix loop.

1. The result

Yu. Il'yshenko [Il] and J. Ecalle [E] proved that an individual planar polynomial vector field can have only finite number of limit cycles.

On the other hand multi-dimensional vector fields with chaotic dynamics have infinite number of periodic trajectories. The Lorentz system [MiMr] and the Duffing system [GuHo] provide best known examples. In the chaotic systems the periodic orbits are usually encoded by periodic sequences in a suitable symbolic dynamical system. This encoding is proved using topological methods (like the Lefschetz-Coneley index or Smale's horseshoe). This means that:

1. The periods of the periodic trajectories tend to infinity in rather irregular way.
2. The 1-cycles represented by different periodic trajectories have different "topology" i.e. they are linked between themselves.

In particular, these cycles do not form a continuous family (so called center).

In Main Theorem below we give an example of polynomial 4-dimensional differential system, with infinite number of periodic solutions $\gamma_1, \gamma_2, \ldots$ such that

- the periods of γ_j grow monotonically with j;
- the corresponding 1-cycles have the same "topology"; they are concentric cycles on an embedded invariant 2-dimensional disc of class C^1;
- the γ_j are isolated (they are limit cycles).

To construct the example we begin with the Hamiltonian planar system

$$\dot{x} = X_H = (H_{x_2}, -H_{x_1}), \quad (x_1, x_2) \in \mathbb{R}^2_x, \qquad H = x_1^3 - 3x_1 - x_2^2 + 2 \tag{1.1}$$

and the 2-dimensional linear system

$$\dot{y} = ay, \qquad y = y_1 + iy_2 \in \mathbb{C} \equiv \mathbb{R}^2_y, \tag{1.2}$$

where $a = -\rho + i\omega$. Later we put $\rho = \omega = \sqrt{3}$.

The Hamiltonian function from (1.1) is elliptic with the critical points $x = (-1, 0)$ (center) and $x = (1, 0)$ (saddle). The phase portrait of the field X_H is shown on Figure 1.1

Date: February 2, 2008.

2000 *Mathematics Subject Classification.* 34C07, 34C08.

This research was supported by the KBN Grant No 2 P03A 015 29.



FIGURE 1.1. Phase portrait of the Hamiltonian vector field X_H with ovals γ_n generating limit cycles

We consider the following coupling of the system (1.1) and (1.2)

$$\begin{cases} \dot{x} = X_H + \mathrm{Re}(\overline{\kappa}\, y)\, e_2 \\ \dot{y} = a\, y + \varepsilon H^4(x)\; (1 - x_1), \end{cases} \tag{1.3}$$

where $\varepsilon > 0$ is a small parameter, $e_2 = (0,1)$ is a versor in $\mathbb{R}^2_x$ and $\kappa \in \mathbb{C}$.

Theorem 1.1 (Main)**.** *Let* $\rho = \omega = \sqrt{3}$ *and*

$$\kappa = 4\sqrt{3}i + \frac{(3-3i)\sqrt{6}}{\sqrt{\pi}}(1 + 2i - \psi'(\tfrac{1-i}{2})), \tag{1.4}$$

where $\psi(z)$*, the Euler Psi-function, is the logarithmic derivative of the Euler Gamma-function* $\psi = \frac{\Gamma'}{\Gamma}$*.*

Then there exists an $\varepsilon_0 > 0$ *such that for any* $0 < \varepsilon < \varepsilon_0$ *the system (1.3) has a sequence of limit cycles* γ_n*,* $n = 1, 2, \ldots$ *which accumulate at the separatrix loop*

$$\gamma_0 = \{(x,y) : \quad y = 0,\ H(x) = 0,\ x_1 \leq 1\}$$

of the singular point $(x = (1,0),\ y = 0)$ *and lie on an invariant surface* $y = \varepsilon G(x, \varepsilon)$ *of class* C^1*.*

Remark 1.2. The approximated numerical value of κ in formula (1.4) is

$$\kappa \approx -0.56 + 4.57i.$$

Systems of the form

$$\begin{cases} \dot{x} = X_H + F(x)y + \varepsilon G(x) + \ldots \\ \dot{y} = A(x)y + \varepsilon b(x) + \ldots, \end{cases} \tag{1.5}$$

$x \in \mathbb{R}^2$, $y \in \mathbb{R}^\nu$, i.e. like (1.3), appear in the so-called multidimensional generalization of the weakened 16-th Hilbert problem (see [Bo, BZ1, BZ2, LZ]). Before perturbation, i.e. for $\varepsilon = 0$, we have the invariant plane $y = 0$ with the Hamiltonian vector field $X_H(x)$. The ovals $H(x) = h$ form a 1-parameter family of its periodic trajectories. One asks how many of these trajectories survive the perturbation. In the 2-dimensional case ($\nu = 0$) the linearization of the problem leads to the problem of real zeroes of an Abelian integral $I(h) = \int_{H(x)=h} \omega$; it is called the weakened 16-th Hilbert problem (see [AI, Il]).

If $\nu \geq 1$, then the corresponding Pontryagin-Melnikov integrals (see [M, P]), denoted $J(h)$, were found in [LZ] and [BZ1]. We call them the generalized Abelian integrals.

The Abelian integrals $I(h)$ satisfy ODEs of the Fuchs type and have regular singularities with real spectrum (see [Yak]). Due to this, S. Yakovenko and others have found some effective estimations for the number of zeroes of $I(h)$. However, the generalized Abelian integrals do not satisfy any simple differential equation (see [Bo]) and sometimes have irregular singularities (e.g. at $h = \infty$). Moreover, even if the singularities are regular, then their spectra can be non-real.

Namely, the non-reality of the spectrum of $J(h)$ at the singularity $h = 0$ is responsible for accumulation of zeroes of J. Below we find the asymptotics

$$J(h) \sim C\, h^{9/2}\, \sin(\log \sqrt{h}), \qquad h \to 0^+.$$

It turns out that the zeroes $h_n \to 0^+$ of J correspond to limit cycles γ_n of the system (1.3); the cycle γ_n bifurcates form the oval $H^{-1}(h_n)$ (see Figure 1.1).

Therefore the system (1.3) can be treated as a counterexample to the multidimensional weakened Hilbert's problem.

The remaining parts of the paper are devoted to the proof of Main Theorem. In Section 2.1 we investigate the generalized Abelian integral and its zeroes. In Section 2.2 we perform estimates needed for existence of genuine limit cycles.

2. Proof of the Main Theorem

2.1. **Generalized Abelian Integral.**

The generalized Abelian integral is defined in two steps. Firstly one solves the so-called *normal variation equation*

$$X_H(g) = ag + (1 - x_1). \tag{2.1}$$

Its solution $x \mapsto g(x) \in \mathbb{C}$ appears in the first (linear in ε) approximation of the invariant surface (see the next section for more details)

$$y = \varepsilon\, H^4(x)\, g(x) + O(\varepsilon^2).$$

We consider (2.1) only in the basin $D = \{x :\ H(x) \geq 0,\, x_1 \leq 1\}$ of the center $x = (-1, 0)$, filled by the periodic solutions $\gamma_h(t) = \gamma(t) \subset \{H^{-1}(h)\}$, each of period

$$T_\gamma(h) = \int_{\gamma_h} -\frac{\mathrm{d}x_1}{2x_2} = \int_{\gamma_h} \mathrm{d}t. \tag{2.2}$$

We assume that the Hamiltonian time is chosen in such a way that for $0 < h < 4$ $x(0) = (x_1^{(1)}, 0)$, where $x_1^{(1)}, x_1^{(2)}, x_1^{(3)}$ are roots of the equation $H(x_1, 0) - h = 0$ (see Figure 2.1). When restricted to γ_h, the equation (2.1) is treated as the ODE $\dot{g} = ag + (1 - x_1)$ with periodic boundary condition. Its unique solution is given in the integral form

$$g(t, h) = (e^{-aT_\gamma} - 1)^{-1} \int_t^{t+T_\gamma} e^{a(t-s)} (1 - x_1)(s, h)\, \mathrm{d}s. \tag{2.3}$$

Substituting the invariant surface equation (see Section 2.2) $y = \varepsilon H^4 g + O(\varepsilon^2)$ into the right hand side of $\dot{x}$ from (1.3) we get the following perturbation of planar Hamiltonian system

$$\begin{cases} \dot{x}_1 = -2x_2, \\ \dot{x}_2 = 3(1 - x_1^2) + \varepsilon\, H^4\, \mathrm{Re}(\overline{\kappa}\, g) + O(\varepsilon^2). \end{cases} \tag{2.4}$$

The generating function for limit cycles is given by the integral

$$J(h) = h^4 \int_{\gamma_h} \mathrm{Re}\big(\overline{\kappa}\, g(x)\big)\, \mathrm{d}x_1. \tag{2.5}$$



FIGURE 2.1. The basin D filled with ovals $\gamma_h \subset H^{-1}(h)$.

Let us denote the "basic" generalized Abelian integral (see [BZ1]) by

$$\Psi_\gamma(h) = \int_{\gamma_h} g(t)(1-x_1)(t)\mathrm{d}t = \\ = (e^{-aT\gamma}-1)^{-1}\int_0^{T_\gamma}\mathrm{d}t\int_t^{t+T_\gamma}\mathrm{d}s\; e^{a(t-s)}(1-x_1)(s)(1-x_1)(t). \tag{2.6}$$

It is related to the generating function via the following

Lemma 2.1. *We have*

$$J(h) = h^4\,\mathrm{Re}\Big[\overline{\kappa}\,\Big(a\Psi_\gamma + 2\int_{\gamma_h}(1-x_1)\,\mathrm{d}t\Big)\Big].$$

Proof. In this proof we denote by dot, $\dot f = \frac{\mathrm{d}}{\mathrm{d}t}f = X_H(f)$, the differential with respect to the Hamiltonian time t. We have

$$J(h) = -h^4\int_\gamma \mathrm{Re}(\overline{\kappa}\,g)\tfrac{\mathrm{d}}{\mathrm{d}t}(1-x_1)\,\mathrm{d}t = h^4\,\mathrm{Re}\Big(\overline{\kappa}\int_\gamma \dot g(1-x_1)\,\mathrm{d}t\Big).$$

Next, $\dot g = a\,g + (1-x_1)$ gives

$$J(h) = h^4\,\mathrm{Re}\Big[\overline{\kappa}\,\Big(a\Psi_\gamma + \int_\gamma (1-x_1)^2\,\mathrm{d}t\Big)\Big] = h^4\,\mathrm{Re}\Big[\overline{\kappa}\Big(a\Psi_\gamma + \int_\gamma (2-2x_1-\tfrac{1}{3}\dot x_2)\,\mathrm{d}t\Big)\Big] = \\ = h^4\,\mathrm{Re}\Big[\overline{\kappa}\Big(a\Psi_\gamma + 2\int_{\gamma_h}(1-x_1)\,\mathrm{d}t\Big)\Big].$$

□

Our next aim is to determine the leading terms in the asymptotic expansion as $h \to 0^+$ of the integrals $\int_\gamma (1-x_1)(t)\mathrm{d}t$, T_γ and Ψ_γ. We begin with the Abelian integrals. It is known [Z2] that these integrals extends to multivalued holomorphic functions with logarithmic singularities. We shall need explicit form of the leading terms.

Lemma 2.2. *There exists an open neighborhood* $0 \in U \subset \mathbb{C}$ *in the complex domain and holomorphic functions* $\eta_0, \eta_1, \zeta_0, \zeta_1 \in \Omega(U)$ *such that*

$$T_\gamma = \eta_0(h) + \zeta_0(h)\, \log h = -\tfrac{1}{2\sqrt{3}} \log h + \tfrac{\sqrt{3}}{2} \log 12 + O(h \log h), \tag{2.7}$$

$$\int_\gamma (1 - x_1) \mathrm{d}t = \eta_1(h) + \zeta_1(h)\, \log h = 2\sqrt{3} + O(h \log h). \tag{2.8}$$

Proof. We consider the pair of basis elliptic Abelian integrals

$$I_0(h) = T_\gamma = \int_\gamma \frac{-\mathrm{d}x_1}{2x_2}, \qquad I_1(h) = \int_\gamma \frac{-x_1 \mathrm{d}x_1}{2x_2}.$$

Note that $\int_\gamma (1 - x_1)\mathrm{d}t = I_0 - I_1$ and that $I_0 = \frac{\mathrm{d}}{\mathrm{d}h}\Big(\text{area of } \{H > h\}\Big)$.

These functions (I_0, I_1) satisfy the Picard-Fuchs equations

$$\begin{aligned} 6h(h-4)I_0' &= -(h-2)I_0 - 2I_1 \\ 6h(h-4)I_1' &= 2I_0 + (h-2)I_1. \end{aligned} \tag{2.9}$$

The other, independent solution to this system is the pair (K_0, K_1), where

$$K_0(h) = \int_{\delta_h} \frac{-\mathrm{d}x_1}{2x_2}, \qquad K_1(h) = \int_{\delta_h} \frac{-x_1 \mathrm{d}x_1}{2x_2}$$

are integrals along another cycle δ_h in the complex curve $E_h = \{H(x) = h\} \subset \mathbb{C}^2$. If $h \in (0, 4)$ then the polynomial $x_1^3 - 3x_1 + 2 - h$ has three real roots $x_h^{(1)} < x_h^{(2)} < x_h^{(3)}$ (see Figure 2.1). The the cycle γ_h (respectively δ_h) is represented as the lift to the Riemann surface E_h of loops in the complex x_1-plane surrounding the roots $x_h^{(1)}$ and $x_h^{(2)}$ (respectively $x_h^{(2)}$ and $x_h^{(3)}$). Note the following integral formulas for $T_\gamma(h)$:

$$T_\gamma = \int_{x_h^{(1)}}^{x_h^{(2)}} \frac{\mathrm{d}x_1}{\sqrt{x_1^3 - 3x_1 + 2 - h}} = \int_{x_h^{(3)}}^{\infty} \frac{\mathrm{d}x_1}{\sqrt{x_1^3 - 3x_1 + 2 - h}}, \qquad h \in (0, 4). \tag{2.10}$$

The second equality corresponds to unobstructed deformation of integration contour γ_h to loop surrounding $x_h^{(3)}$ and ∞.

The system (2.9) has resonant singular point $h = 0$. Any its solution is either analytic near $h = 0$ (it is (K_1, K_2)) or it has the form like I_0, I_1:

$$\begin{aligned} I_0(h) &= (a_0 + a_1 h + \ldots) + \tfrac{1}{2\pi i} K_0 \, \log h, \\ I_1(h) &= (b_0 + b_1 h + \ldots) + \tfrac{1}{2\pi i} K_1 \, \log h. \end{aligned} \tag{2.11}$$

This representation follows from the Picard-Lefschetz formula

$$\gamma_h \longrightarrow \gamma_h \cdot \delta_h, \qquad \delta_h \longrightarrow \delta_h, \tag{2.12}$$

which describes the monodromy transformations of the generators of $\pi_1(E_h, *)$, as h surrounds the critical value 0; here $*$ denotes a basepoint.

We need to calculate the expansions of I_0, I_1. As we shall see, it is enough to calculate $K_0(0)$ and a_0; all other coefficients follows from the system (2.9) and can be recursively determined. Indeed, to compensate terms with $\log h$ in (2.9) we must have

$$K_1(0) = K_0(0). \tag{2.13}$$

Terms with h^0 give

$$b_0 = a_0 + \tfrac{12}{2\pi i} K_0(0). \tag{2.14}$$

It can be continued further.

To determine $K_0(0)$ and a_0 simultaneously, we make a coordinate change $u = (x_1-1)/(x_h^{(3)}-1)$ in the integral (2.10); we denote also $p = (x_h^{(3)}-1)$. Since $p = \sqrt{h/3}+O(h)$ as $h\to 0^+$, the following integral

$$\int_1^\infty \mathrm{d}u\Big[\frac{1}{\sqrt{u^2(3+pu)-h/p^2}} - \frac{1}{\sqrt{u^2(3+pu)}} - \frac{1}{\sqrt{3u^2-3}} + \frac{1}{\sqrt{3}}\Big] \xrightarrow{h\to 0^+} 0.$$

We calculate

$$\int_1^\infty \mathrm{d}u\Big[\frac{1}{\sqrt{3u^2-3}} - \frac{1}{\sqrt{3}}\Big] = \frac{\log 2}{\sqrt{3}},$$

$$\int_1^\infty \mathrm{d}u\Big[\frac{1}{u\sqrt{3+pu}}\Big] = \tfrac{2}{\sqrt{3}}\log\Big(\tfrac{2\sqrt{3}}{\sqrt{p}}+o(1)\Big) = -\tfrac{1}{2\sqrt{3}}\log h + \frac{\log(12\sqrt{3})}{\sqrt{3}} + o(h^{1/2}).$$

Thus $a_0 = \frac{\sqrt{3}}{2}\log 12$, $K_0(0) = -\frac{2\pi i}{2\sqrt{3}}$. Substituting these values to the relations (2.13), (2.14) and using the expansion (2.11) we get the leading terms of the expansions as in formulas (2.7) and (2.8).

□

Let us pass to expansion of Ψ_γ.

Proposition 2.3. *Let $-2\sqrt{3} < \mathrm{Re}(a) < 0$. There exists an open neighborhood $0\in U\subset\mathbb{C}$ in the complex domain and holomorphic functions $\varphi_1,\varphi_2,\varphi_3$ such that*

(2.15)

$$\Psi_\gamma(z) = \varphi_1(h)+\varphi_2(h)\,\log h+\varphi_3(h)\cdot\Big(e^{-aT_\gamma}-1\Big)^{-1} = C_0 + C_1 h^{-a/2\sqrt{3}}+\ldots,$$

where

$$C_1 = \frac{(\pi a)^2}{\sin^2(\pi a/2\sqrt{3})}, \tag{2.16}$$

$$C_0 = \frac{3\sqrt{2}}{\sqrt{\pi}}\Big(-1+2w+2w^2\psi'(-w)\Big), \qquad w = \tfrac{a}{2\sqrt{3}}. \tag{2.17}$$

Remark 2.4. One can easily observe that the value (1.4) of κ satisfies the relation

$$\kappa = i(4\sqrt{3}+aC_0). \tag{2.18}$$

It is chosen in a way to annihilate the leading term ($\sim h^4$) of $J(h)$ and to reveal the term with the infinite sequence of zeroes – see the Corollary 2.5 and its proof below.

Corollary 2.5. *Providing the values of parameters (ρ,ω,κ) as in Theorem Main, the integral $J(h)$ (2.5) has a sequence h_n, $n=1,2,\ldots$ of simple zeroes accumulating at $h=0$.*

Proof. We calculate the leading term of the expansion of $J(h)$ using Lemma 2.1, Lemma 2.2 and Remark 2.4:

$$\begin{aligned} J(h) &= h^4\mathrm{Re}\Big[\overline{\kappa}\left(aC_0+aC_1h^{1/2-i/2}+4\sqrt{3}+o(h^{3/4})\right)\Big] = \\ &= h^4\mathrm{Re}\Big[\overline{\kappa}\left(aC_0+4\sqrt{3}\right)\Big] + h^4\mathrm{Re}\Big[\overline{\kappa}\,a\,C_1h^{1/2-i/2}\Big] + o(h^{4+3/4}) = \\ &= R\,h^{4+1/2}\cos(\log\sqrt{h}-\alpha_0)+o(h^{4+3/4}), \end{aligned}$$

where $R = |\overline{\kappa}\,a\,C_1|$, $\quad\alpha_0 = \arg(\overline{\kappa}\,a\,C_1)$. Analogously we get

$$J'(h) = R_1\,h^{3+1/2}\cos(\log\sqrt{h}-\alpha_1)+o(h^{3+3/4}).$$

Thus, by Implicit Function Theorem the zeroes (h_n) of $J(h)$ approximate the simple zeroes $h_n^{(0)}$ of the function $\cos(\log\sqrt{h}-\alpha_0)$.

$\square$

The remaining part of this section is devoted to the proof of Proposition 2.3. It goes in two steps. In first one we show that the function

$$\Psi_\gamma(h) \longrightarrow C_0$$

as $h \to 0^+$ and so is bounded.

In the second step we determine the monodromy of the generalized Abelian integral $\Psi_\gamma(h)$ as h surrounds $h=0$. We know that then γ changes to $\mathcal{M}\mathrm{on}_0\gamma = \gamma\cdot\delta$ and $\mathcal{M}\mathrm{on}_0\delta = \delta$. We would like to express $\Psi_{\gamma\cdot\delta}$ in simple terms, in order to determine the singularity of $\Psi_\gamma(h)$ at $h=0$. Rather complicated formulas for $\Psi_{\gamma\cdot\delta}$ are given in [BZ1] and [BZ2]. In [Bo] these formulas were simplified using certain upper triangle representation ρ of the fundamental group $\pi_1(E_h,*)$. We recall this construction below.

We denote

$$\begin{aligned}
\Psi_\gamma(h) &= (e^{-aT\gamma}-1)^{-1}\int_0^{T_\gamma}\mathrm{d}t\int_t^{t+T_\gamma}\mathrm{d}s\; e^{a(t-s)}(1-x_1)(s)\,(1-x_1)(t),\\
\lambda_\gamma(h) &= e^{-aT_\gamma/2},\\
\phi_\gamma(h) &= \lambda_\gamma\int_0^{T_\gamma}\mathrm{d}t\int_0^t\mathrm{d}s\;(1-x_1)(t)\,(1-x_1)(s)\cdot e^{a(t-s)} =\\
&= \lambda_\gamma\int_{-T_\gamma/2}^{T_\gamma/2}\mathrm{d}t\int_{-T_\gamma/2}^t\mathrm{d}s\;(1-x_1)(t+\tfrac{T_\gamma}{2})\,(1-x_1)(s+\tfrac{T_\gamma}{2})\,e^{a(t-s)},\\
\theta_\gamma^+ &= \lambda_\gamma\int_0^{T_\gamma}\mathrm{d}t\;(1-x_1)(t)\,e^{at} = \int_{-T_\gamma/2}^{T_\gamma/2}\mathrm{d}t\;(1-x_1)(t+\tfrac{T_\gamma}{2})\,e^{at},\\
\theta_\gamma^- &= \lambda_\gamma^{-1}\int_0^{T_\gamma}\mathrm{d}t\;(1-x_1)(t)\,e^{-at} = \int_{-T_\gamma/2}^{T_\gamma/2}\mathrm{d}t\;(1-x_1)(t+\tfrac{T_\gamma}{2})\,e^{-at},
\end{aligned} \tag{2.19}$$

Here the subscript γ underlines dependence of the above functions on the loop $\gamma=\gamma_h$.

We introduce the following space of triangular matrices

$$\mathbb{T}\colon = \left\{ \begin{pmatrix} \lambda & \theta^- & \phi\\ 0 & \lambda^{-1} & \theta^+\\ 0 & 0 & \lambda \end{pmatrix}, \quad \lambda\in\mathbb{C}^*, \quad \theta^+,\theta^-,\phi\in\mathbb{C} \right\}; \tag{2.20}$$

it forms a group. For $W\in\mathbb{T}$ we denote

$$|W| = \det W = \lambda.$$

Existence of a 2-dimensional Jordan cell is measured in the following formula

$$\frac{(W-|W|)(W-1/|W|)}{|W|^2-1} = \psi(W)\begin{pmatrix} 0&0&1\\ 0&0&0\\ 0&0&0 \end{pmatrix}; \tag{2.21}$$

explicitly we have

$$\psi(W) = \frac{\theta^+\theta^-}{\lambda^2-1} + \frac{\phi}{\lambda}. \tag{2.22}$$

Theorem 2.6 ([Bo]). *The map* $\rho : \pi_1(E_h,*) \to \mathbb{T}$,

$$\rho(\gamma) = \begin{pmatrix} \lambda_\gamma & \theta_\gamma^- & \phi_\gamma\\ 0 & \lambda_\gamma^{-1} & \theta_\gamma^+\\ 0 & 0 & \lambda_\gamma \end{pmatrix}, \tag{2.23}$$

where $\lambda_\gamma, \theta_\gamma^\pm, \phi_\gamma$ *are defined in (2.19), defines a representation of the fundamental group of* E_h*. Moreover, we have*

$$\Psi_\gamma = \psi \circ \rho\,(h). \tag{2.24}$$

Sketch of the proof. We have $\Psi_\gamma = (\lambda_\gamma^2 - 1)^{-1} \iint e^{a(t-s)}(1-x)(t)\,(1-x)(s)$, where the integration domain is $\Sigma = \{(t,s):\ 0 \le t \le T_\gamma,\ t \le s \le T_\gamma + t\}$ (see (2.19)). We divide Σ into two "triangles" $\triangle_1 = \{0 \le t \le T_\gamma,\ t \le s \le T_\gamma\}$ and $\triangle_2 = \{0 \le t \le T_\gamma,\ T_\gamma \le s \le T_\gamma + t\} = \triangle_0 + (0, T_\gamma)$, where $\triangle_0 = \{0 \le t \le T_\gamma,\ 0 \le s \le t\}$. We have $\iint_{\triangle_1+\triangle_2}(\cdot) = \iint_{\triangle_1+\triangle_0}(\cdot) + \iint_{\triangle_2-\triangle_0}(\cdot)$, where $\iint_{\triangle_1+\triangle_0}(\cdot) = \theta_\gamma^+\,\theta_\gamma^-$ and $\iint_{\triangle_2-\triangle_0}(\cdot) = (\lambda_\gamma^2 - 1)\lambda_\gamma^{-1}\phi_\gamma$. Now the formula (2.24) follows from (2.22).

The property $\rho(\gamma\cdot\delta) = \rho(\gamma)\ \rho(\delta),\ \gamma, \delta \in \pi_1(E_h, *)$ is proved analogously. We divide the line integrals in $\theta_{\gamma\delta}^\pm$ and the surface integral in $\phi_{\gamma\delta}$ into parts where t or s lies in γ or in δ. We use also $\lambda_{\gamma\delta} = \lambda_\gamma\lambda_\delta$.

□

Proposition 2.7. *Let* $\xi(t) = (1-x_1)(t)$ *with the initial value* $\xi(0) = 1 - x_1^{(1)}$ *(see Figure 2.1). We have the following integral formula for the generalized Abelian integral*

$$\Psi_\gamma(h) = (e^{-aT_\gamma} - 1)^{-1}\Big(\int_{-T_\gamma/2}^{T_\gamma/2} \xi(t)e^{at}\mathrm{d}t\Big)^2 + \int_{-T_\gamma/2}^{T_\gamma/2}\mathrm{d}t\int_{-T_\gamma/2}^{t}\mathrm{d}s\ \xi(s)\xi(t)e^{a(t-s)}. \tag{2.25}$$

As $h \to 0^+$ *these integrals have finite limits:*

$$\begin{aligned} &\int_{-T_\gamma/2}^{T_\gamma/2} \xi(t)e^{at}\mathrm{d}t \longrightarrow \frac{\pi a}{\sin(\pi a/2\sqrt{3})} = \sqrt{C_1}, \\ &\int_{-T_\gamma/2}^{T_\gamma/2}\mathrm{d}t\int_{-T_\gamma/2}^{t}\mathrm{d}s\ \xi(s)\xi(t)e^{a(t-s)} \longrightarrow C_0, \end{aligned} \tag{2.26}$$

where C_0, C_1 *are as defined in Proposition 2.3.*

Proof. The value of generalized Abelian integral Ψ_γ does not depend on the "shift" of parametrization (e.g. $t \mapsto t + \frac{T_\gamma}{2}$), but values of integrals ϕ_γ and $\theta_\gamma^\pm$ depend. We choose Hamiltonian time parameter in such a way that $x_1(0) = x_1^{(2)}$, where $x_h^{(1)} < x_h^{(2)} < x_h^{(3)}$ are real roots of the polynomial $x_1^3 - 3x_1 + 2 = h$ (see Figure 2.1). Thus

$$(1-x_1)(t + T_\gamma/2) = \xi(t)$$

and so, using formula (2.22) and formulas (2.19), we get formula (2.25).

To determine the asymptotic expansion we notice that the singular curve $E_0 = \{H(x) = 0\} = \{x_2^2 = (x_1-1)^2(x_1+2)\}$ is rational. The Hamiltonian parametrization of the limit loop γ_0 can be explicitly calculated:

$$\xi(t) \longrightarrow \xi_0(t) = \frac{3}{\cosh^2(\sqrt{3}t)}, \qquad -\infty < t < \infty \tag{2.27}$$

as $h \to 0^+$. Recall that $T_\gamma(0) = \infty$. Substituting these values to integrals in (2.25) we get the following limits

$$\int_{-T_\gamma/2}^{T_\gamma/2} \xi(t)e^{at}\mathrm{d}t \longrightarrow \int_{-\infty}^{\infty} \frac{3}{\cosh^2(\sqrt{3}t)}e^{it(a/i)}\mathrm{d}t = \sqrt{2\pi}\mathcal{F}(\xi_0)(a/i)$$

where $\mathcal{F}$ denotes the Fourier transform. Since $\mathcal{F}(\frac{1}{\cosh^2})(k) = \sqrt{\frac{\pi}{2}}\frac{k}{\sinh(k\pi/2)}$ (see [GR], Integral 3.982.1 for example) we find the value

$$\frac{i\pi a}{\sinh(i\pi a/2\sqrt{3})} = \frac{\pi a}{\sin(\pi a/2\sqrt{3})} = \sqrt{C_1}.$$

To determine the limit of the second integral

$$\int_{-T_\gamma/2}^{T_\gamma/2} \mathrm{d}t \int_{-T_\gamma/2}^{t} \mathrm{d}s\ \xi(t)\xi(s)e^{a(t-s)} \longrightarrow \int_{-\infty}^{\infty} \mathrm{d}t \int_{-\infty}^{t} \mathrm{d}s\ \xi_0(t)\xi_0(s)e^{a(t-s)}$$

we substitute $u = t - s$, use the symmetry $\xi_0(-t) = \xi_0(t)$ and the Parsival identity; then we get

$$\int_{-\infty}^{\infty} \mathrm{d}t \int_{0}^{\infty} \mathrm{d}u\ \xi_0(t)\xi_0(u-t)e^{a(u)} = \int_{-\infty}^{\infty} \mathrm{d}k \Big(\mathcal{F}(\xi_0)(k)\Big)^2 \mathcal{F}\Big(e^{au}\chi_{[0,\infty)}(u)\Big)(k) =$$

$$= \frac{3\sqrt{2}i}{\pi^{3/2}} \int_{-\infty}^{\infty} \frac{k^2}{\sinh^2 k} \frac{\mathrm{d}k}{k - \pi i(a/2\sqrt{3})}.$$

To evaluate the latter integral, which has the form

$$F(w) = \int_{-\infty}^{\infty} \frac{z^2}{\sinh^2 z} \frac{\mathrm{d}z}{z - \pi i w}, \qquad \mathrm{Re} w < 0, \tag{2.28}$$

we must use the logarithmic derivative of the Euler Γ function, i.e. the function $\psi = (\log \Gamma)' = \frac{\Gamma'}{\Gamma}$.

Integrating by parts we obtain

$$F(w) = \lim_{R\to+\infty} \int_{-R}^{-R^{-1}} + \int_{R^{-1}}^{R} (\mathrm{sgn} z - \coth z)' \frac{z^2}{z - \pi i w} \mathrm{d}z =$$

$$= -2\pi i w + \pi^2 w^2 \lim_{R\to+\infty} \int_{-R}^{-R^{-1}} + \int_{R^{-1}}^{R} \frac{\coth z}{(z - \pi i w)^2} \mathrm{d}z.$$

Next, we integrate the function $\frac{\coth z}{(z-\pi i w)^2}$ along the contour consisting of the segment $[-R, -R^{-1}]$ followed by the semicircle $R^{-1}e^{i\varphi}$, $\varphi \in [\pi, 2\pi]$ and segments: $[R^{-1}, R]$, $[R, R + i(N + \frac{1}{2})\pi]$, $[R + i(N + \frac{1}{2})\pi, -R + i(N + \frac{1}{2})\pi]$, $[-R, -R + i(N + \frac{1}{2})\pi]$, where $N \in \mathbb{N}$. Using the residue formula and passing to the limit $R, N \to \infty$ we deduce

$$\lim_{R\to+\infty} \int_{-R}^{-R^{-1}} + \int_{R^{-1}}^{R} \frac{\coth z}{(z - \pi i w)^2} \mathrm{d}z + \frac{\pi i}{(\pi i w)^2} = 2\pi i \sum_{n=0}^{\infty} \mathrm{Res}_{i\pi n}\Big(\frac{\coth z}{(z - \pi i w)^2}\Big) =$$

$$= -\frac{2i}{\pi} \sum_{n=0}^{\infty} \frac{1}{(n-w)^2} = -\frac{2i}{\pi}\, \psi(-w);$$

in the identification of the latter sum we used [GR], formula 8.363.8. Finally we have

$$F(w) = \pi i\Big(1 - 2w - 2w^2\psi'(-w)\Big) \qquad \text{for} \quad \mathrm{Re}(w) < 0,$$

and so the second of the limits (2.26) follows.

□

Now we investigate the monodromy properties of the generalized Abelian integral Ψ_γ. We shall need the following

Lemma 2.8 ([Bo]). *For $W, W' \in \mathbb{T}$ we have*

$$\psi(W \cdot W') = \psi(W) + \psi(W') + \frac{|W|^2\,|W'|^2}{(|W|^2 - 1)(|W'|^2 - 1)(|W|^2\,|W'|^2 - 1)} \widetilde{\psi}([W, W']),$$

where $[W, W'] = W\, W'\, W^{-1}\, (W')^{-1}$ is the commutant and

$$\widetilde{\psi}(W) = (|W|^2 - 1)\ \psi(W);$$

(for $|W| = 1$ we have $\widetilde{\psi}(W) = \theta^+\,\theta^-$ in terms of (2.20)).

Proof. The proof relies on direct calculations.

□

Corollary 2.9. *The function $\Psi_\gamma(h)$ near $h=0$ has the following form*

$$\Psi_\gamma(h)=\varphi_1(h)+\tfrac{1}{2\pi i}\Psi_\delta(h)\log h-\frac{\lambda_\delta^2(h)}{(\lambda_\delta^2(h)-1)^2}\widetilde{\Psi}_{[\gamma,\delta]}(h)\cdot\frac{1}{\lambda_\gamma(h)-1}, \tag{2.29}$$

where δ is the second cycle in $\pi_1(E_h,*)$ (see the proof of Lemma 2.2), $\widetilde{\Psi}_{[\gamma,\delta]}(h)=\widetilde{\psi}(\rho([\gamma,\delta])$. The functions φ_1, Ψ_δ and $\widetilde{\Psi}_{[\gamma,\delta]}$ are holomorphic near $h=0$.

Remark 2.10. One can prove that the function $\widetilde{\Psi}_{[\gamma,\delta]}$ is constant

$$\widetilde{\Psi}_{[\gamma,\delta]}(h)=(2\pi a)^2.$$

Indeed, since the contour $[\gamma,\delta]$ is monodromy invariant (see proof of Corollary 2.9 below), the function $\widetilde{\Psi}_{[\gamma,\delta]}$ is meromorphic on whole $\mathbb{C}$ with possible poles in $h=0,4$. We know, by Proposition 2.7, that it is bounded as $h\to 0$. Similarly one shows that it is bounded as $h\to 4$. These calculations are analogous to proof of the first limit in (2.26). We also check that $\widetilde{\Psi}_{[\gamma,\delta]}$ is bounded as $h\to\infty$. Thus this function has to be constant; its value we calculate by passing to the limit $h\to 0$ and comparing respective terms in (2.25) and (2.29).

Proof of Corollary 2.9. The Picard-Lefschetz formula (2.12), Theorem 2.6 and Lemma 2.8 imply that

$$\mathcal{M}\mathrm{on}_0\Psi_\gamma=\psi(\rho(\gamma)\cdot\rho(\delta))=\Psi_\gamma+\Psi_\delta+\frac{\lambda_\gamma^2\lambda_\delta^2}{(\lambda_\gamma^2-1)\,(\lambda_\delta^2-1)\,(\lambda_\gamma^2\lambda_\delta^2-1)}\widetilde{\Psi}_{[\gamma,\delta]} \tag{2.30}$$

and $\mathcal{M}\mathrm{on}_0\Psi_\delta=\Psi_\delta$. Next the following monodromy relations follows the Picard-Lefschetz formula (2.12)

$$\mathcal{M}\mathrm{on}_0\lambda_\gamma=\lambda_\gamma\,\lambda_\delta,\qquad \mathcal{M}\mathrm{on}\lambda_\delta=\lambda_\delta,$$
$$\mathcal{M}\mathrm{on}_0[\gamma,\delta]=(\gamma\delta)\cdot\delta\cdot(\gamma\delta)^{-1}\cdot\delta^{-1}=[\gamma,\delta].$$

Therefore Ψ_δ, $\widetilde{\Psi}_{[\gamma,\delta]}$ and λ_δ are locally single-valued functions of h. Since they are bounded (see Proposition 2.7), they must be holomorphic. Now

$$\mathcal{M}\mathrm{on}_0\Big(\tfrac{1}{2\pi i}\Psi_\delta\,\log h\Big)=\tfrac{1}{2\pi i}\Psi_\delta\,\log h+\Psi_\delta,$$

$$\begin{aligned}\mathcal{M}\mathrm{on}_0\Big(-\frac{\lambda_\delta^2\,\widetilde{\Psi}_{[\gamma,\delta]}}{(\lambda_\delta^2-1)^2\,(\lambda_\gamma^2-1)}\Big)&=-\frac{\lambda_\delta^2\,\widetilde{\Psi}_{[\gamma,\delta]}}{(\lambda_\delta^2-1)^2\,(\lambda_\gamma^2\lambda_\delta^2-1)}=\\&=\Big(-\frac{\lambda_\delta^2\,\widetilde{\Psi}_{[\gamma,\delta]}}{(\lambda_\delta^2-1)^2\,(\lambda_\gamma^2-1)}\Big)+\frac{\lambda_\gamma^2\lambda_\delta^2\,\widetilde{\Psi}_{[\gamma,\delta]}}{(\lambda_\gamma^2-1)\,(\lambda_\delta^2-1)\,(\lambda_\gamma^2\lambda_\delta^2-1)}.\end{aligned}$$

Therefore the function φ_1 defined by (2.29) is single-valued. Since (λ_γ^2-1) and (λ_δ^2-1) are separated from zero and the function Ψ_γ is bounded (see Proposition 2.7), the function φ_1 is holomorphic.

□

Corollary 2.9 allows to finish the proof of Proposition 2.3. The holomorphic function φ_1 is defined in Corollary and φ_2, φ_3 can be read from (2.29):

$$\varphi_2=\tfrac{1}{2\pi i}\Psi_\delta,$$
$$\varphi_3=-\frac{\lambda_\delta^2\,\widetilde{\Psi}_{[\gamma,\delta]}}{(\lambda_\delta^2-1)^2}=-\Big(\frac{2\pi a\lambda_\delta}{\lambda_\delta^2-1}\Big)^2$$

(the latter equality follows from Remark 2.10).

Since $T_\gamma=-\frac{1}{2\sqrt{3}}\log h+O(1)$, we have the leading term of expansion

$$(\lambda_\gamma^2-1)^{-1}=(e^{-aT_\gamma}-1)^{-1}=h^{-a/2\sqrt{3}}+\ldots$$

□

2.2. **Estimates.** In this subsection we show that the zeroes h_n of the generalized Abelian integral (see Corollary 2.5) generate corresponding limit cycles of the system (1.3), provided ε is sufficiently small.

Recall that the problem of limit cycles of (1.3) is reduced to the problem of limit cycles of the following planar system

$$\dot{x} = X_H(x) + \varepsilon \operatorname{Re}\Big(\overline{\kappa}\, G(x,\varepsilon)\Big) e_2, \tag{2.31}$$

where the function $G(x,\varepsilon)$ is defined via the invariant surface $L_\varepsilon = \{y = \varepsilon G(x,\varepsilon)\}$, which is a graph of $G(\cdot,\varepsilon)$.

At the moment we do not even know whether the invariant surface exists. Indeed, the normal hyperbolicity conditions are *not* satisfied: the eigenvalues in the normal direction are $\lambda_{3,4} = -\sqrt{3} \pm i\sqrt{3}$, whereas the eigenvalues at the saddle point $x = (1,0), y = 0$ in the x-direction are $\pm 2\sqrt{3}$ (compare [BZ1, BZ2, HPS, Ni]). We should do two things:

(1) prove the existence of the invariant surface,
(2) estimate the discrepancy $G(x,\varepsilon) - H^4\, g(x)$, where $g(x)$ is the solution to the normal variation equation given in (2.1).

In both tasks the crucial role is played by the following Lemma. Let us recall the notation related to the elliptic Hamiltonian $H(x) = x_1^3 - 3x_1 - x_2^2 + 2$. The basin $D \subset \mathbb{R}^2$ (see Figure 2.1) is filled with closed orbits of the Hamiltonian vector field X_H.

Lemma 2.11. *Let $U \supset D \times \{0\}$ be an open neighborhood in $\mathbb{R}^2 \times \mathbb{C}$ and V_ε be the following vector field in U*

$$V_\varepsilon \begin{cases} \dot{x} = X_H + H^k \operatorname{Re}(\overline{\kappa}\, y) v_o + Q(x,y;\varepsilon) \\ \dot{y} = ay + B(x,y;\varepsilon), \end{cases} \tag{2.32}$$

where $k \geq 0$, $a = -\sqrt{3} + i\sqrt{3}$, $\kappa \in \mathbb{C}$, $v_0 \in \mathbb{R}^2$ and Q, B are functions of class $C^2(U)$ satisfying the following

$$|Q| \leq Const \cdot \varepsilon |H(x)|^2,$$
$$|B| \leq Const \cdot \varepsilon |H(x)|^2.$$

Then, for sufficiently small ε there exists a unique invariant surface of V_ε:

$$L_\varepsilon = \{(x,y) : \quad x \in D,\ y = \varepsilon\, G(x,\varepsilon)\}.$$

The function $G(\cdot,\varepsilon)$ prolongs by zero to a C^1 function on a neighborhood of D in $\mathbb{R}^2_x$.

Proof. We shall prove that the Poincaré return map associated to vector field V_ε satisfies the normal hyperbolicity condition.

In a neighborhood of the center critical point $(x = (-1,0), y = 0)$ for unperturbed vector field V_0, the system is normally hyperbolic and so the invariant surface exists. In the further proof we shall concentrate on the neighborhood of separatrix $\gamma_0 = \{(x,y) : y = 0, \quad H(x) = 0\}$ (see Figure 2.2).

The non-degenerate critical point $p_0 = ((1,0),0)$, located on this separatrix, is preserved after the perturbation. Let us choose a 3-dimensional hypersurface S transversal to γ_0 and close to the singular point p_0. Let

$$\Sigma_{\alpha,d} = S \cap \{(x,y) : \quad |y|^2 \leq \alpha^2 (H(x))^2, \quad H(x) < d\}$$

be a sector in S with vertex

$$\sigma_0 = \Sigma_{\alpha,d} \cap \gamma_0 = \{(x,0) \in \Sigma_{\alpha,d} : \quad H(x) = 0\}.$$



FIGURE 2.2. The Poincaré return map $\mathcal{P}_\varepsilon$ defined via trajectories of V_ε and its invariant unit I_ε.

Lemma 2.12. *For sufficiently small* $\alpha > 0$, $d_2 > d_1 > 0$ *and* $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$, *the Poincaré return map*

$$\mathcal{P}_\varepsilon : \Sigma_{\alpha,d_1} \to \Sigma_{\alpha,d_2} \tag{2.33}$$

is a diffeomorphism onto the image and prolongs to a map of class C^1 *in point* σ_0.

Now we finish the proof of Lemma 2.11. The unperturbed Poincare map $\mathcal{P}_0$ is the identity on the invariant segment $I_0 = \Sigma_{\alpha,d_1} \cap \{y = 0\}$. Thus $\mathcal{P}_0$ is normally hyperbolic on I_0, since we have strong contraction in the normal direction. In virtue of the Hirsh Pugh Shub Theorem [HPS], for sufficiently small ε there exist the unique invariant embedded interval I_ε close to I_0; it is of class C^1. Considering Hamiltonian as the parameter on I_0, we get

$$I_\varepsilon = \{y = \varepsilon F(h, \varepsilon),\ h \in [0, \delta)\}, \quad F \in C^1([0, \delta) \times (-\varepsilon_0, \varepsilon_0)).$$

The surface S_ε spanned by trajectories of V_ε passing through I_ε is V_ε-invariant, due to the invariance of I_ε under the Poincaré map $\mathcal{P}_\varepsilon$. The form of invariant interval I_ε and the form of vector field V_ε implies that the surface S_ε is graph of a $C^1(D)$ function:

$$L_\varepsilon = \{(x, y) : x \in D, \quad y = \varepsilon G(x, \varepsilon)\}.$$

We prove that G can be extended by zero outside D.

We check that, providing the assumptions of Lemma 2.11 hold, the set

$$\{(x, y) : x \in D, |y|^2 \le R^2 \varepsilon^2 |H(x)|^4\},$$

for R big enough, is invariant for V_ε. Namely, denoting all constants by C, we calculate

$$\begin{aligned} &V_\varepsilon(|y|^2 - R^2\varepsilon^2|H|^4)|_{|y|=R\,\varepsilon\,|H|^2} = \\ &= 2\mathrm{Re}\Big[\overline{y}\ (a\,y + B)\Big] - 4R^2\varepsilon^2 H^3\Big(H^3\mathrm{Re}(\overline{\kappa}\,y\ < \mathrm{d}H, v_0 > + < \mathrm{d}H, Q >)\Big) \\ &\qquad \le 2R^2\,\varepsilon^2\,H^4\Big(-\rho + \tfrac{C}{R} + 2R\,\varepsilon\,C\,|H|^{1+k} + \varepsilon\,C\,|H|\Big). \end{aligned} \tag{2.34}$$

Since the latter expression is ≤ 0 (for sufficiently large R), the considered subset is V_ε-invariant.

Thus

$$|G(x,\varepsilon)| \leq Const \cdot |H(x)|^2$$

and the function $G(x,\varepsilon)$ can be prolonged by zero to a function of class C^1.

□

Proof of Lemma 2.12. By calculations analogous to (2.34) we find that

$$V_\varepsilon(|y|^2-\alpha^2 H^2)|_{|y|=\alpha\,|H|} \leq 2\alpha^2\, H^2\Big(-\rho+\tfrac{\varepsilon\,|H|}{\alpha}+\alpha\, C\,|H|^k + C\varepsilon\,|H|\Big).$$

Thus for sufficiently small α and ε the subset

$$\{(x,y): x\in D, \quad |y|^2 \leq \alpha^2 (H(x))^2\}$$

is V_ε-invariant. Moreover, the separatrix γ_0 is also V_ε-invariant. This proves, that the Poincaré return map defines the dyffeomorphism (2.33) which is of class C^1 outside the border.

Thus it remains to show that $\mathcal{P}_\varepsilon$ can be prolonged to the C^1 map in the point σ_0. We choose an additional, auxiliary, 3-dimensional hypersurface $\widetilde{S}$ transversal to γ_0, close to p_0, which lies "on another side" with respect to the point p_0 (see Figure 2.2). The return map $\mathcal{P}_\varepsilon$ is the composition $\mathcal{P}_\varepsilon = \mathcal{P}^s_\varepsilon \circ \mathcal{P}^r_\varepsilon$ of the correspondence maps

$$\mathcal{P}^s_\varepsilon : S \to \widetilde{S}, \qquad \text{and} \qquad \mathcal{P}^r_\varepsilon : \widetilde{S} \to S,$$

defined by trajectories near the singular point p_0 and trajectories near the regular part of γ_0 respectively. The regular map naturally extends to the C^1 map (even C^2 in fact) as the flow of the non-vanishing, C^2 vector field V_ε. To analyze the singular part $\mathcal{P}^s_\varepsilon$, we use the following theorem of H. Belitskii.

Theorem 2.13 ([Be])**.** *Let* $\Lambda \in \mathrm{End}(\mathbb{R}^n)$ *be a linear endomorphism whose eigenvalues* $(\lambda_1,\ldots,\lambda_n)$ *satisfy*

$$\mathrm{Re}\lambda_i \neq \mathrm{Re}\lambda_j + \mathrm{Re}\lambda_k$$

for all i *and* j,k *such that* $\mathrm{Re}\lambda_j \leq 0 \leq \mathrm{Re}\lambda_k$. *Then any* C^2 *differential system*

$$\frac{\mathrm{d}x}{\mathrm{d}t} = \Lambda x + f(x), \qquad f(0) = 0 = f'(0)$$

is in the neighborhood of 0 C^1*-equivalent to the linearization.*

The eigenvalues of the linearization of our vector field V_ε in p_0 are $\pm 2\sqrt{3}$, $-\sqrt{3}\pm i\sqrt{3}$, so V_ε satisfies the assumptions of the Belitskii theorem. In suitable coordinates (u,v), associated with the linearization of V_ε in the neighborhood of p_0, the correspondence map $\mathcal{P}^s_\varepsilon$ has the form

$$\mathcal{P}^s_\varepsilon(u,v) = (u, C\, u^\beta\, v), \quad u\in\mathbb{R}_+, \quad v\in(\mathbb{C},0), \quad \beta = \tfrac{1}{2}-\tfrac{i}{2}.$$

The restriction to $\Sigma_{\alpha,d}$ corresponds to the restriction to the set $\{|v|^2 \leq \widetilde{\alpha}(u)\,u,\ u\in\mathbb{R}_+,\ \widetilde{\alpha}(0)>0\}$. In such region the map $\mathcal{P}^s_\varepsilon$ is of class C^1 in σ_0; we have $(u,v)(\sigma_0) = (0,0)$ and $(\mathcal{P}^s_\varepsilon)'(0,0) = \left(\begin{smallmatrix}1&0\\0&0\end{smallmatrix}\right)$. Thus, the thesis of the lemma follows.

□

Now we can prove the existence of the invariant surface and estimate the distance to its linear approximation $H^4 g$.

Proposition 2.14. *For sufficiently small* ε, *there exists the invariant surface*

$$L_\varepsilon = \{y = \varepsilon\, G(x,\varepsilon), \quad x\in D\} \tag{2.35}$$

of the system (1.3). The function G *is of class* C^1 *and the distance to the linearization* $(H^4 g)(x) = G(x,0)$ *is bounded by*

$$|G - H^4 g| \leq C|\varepsilon||h|^5, \tag{2.36}$$

$$|G' - (H^4 g)'| \leq C|\varepsilon||h|^4, \tag{2.37}$$

where $G' = G_{,x}$ *is the derivative with respect to* x*.*

Proof. The existence of the invariant surface of class C^1 and the form (2.35) is a direct consequence of Lemma 2.11.

To show the bounds (2.36), (2.37), we make a coordinate change

$$(x,y) \longmapsto (x,z), \qquad z = \Big(y - \varepsilon\, H^4 g(x)\Big)/(\varepsilon\, H^4).$$

The system (1.3) takes the form

$$\begin{cases} \dot{x} =X_H + H^4\,\mathrm{Re}(\overline{\kappa}\, z)\,\varepsilon\, e_2 + \varepsilon\, H^4\,\mathrm{Re}(\overline{\kappa}\, g)\, e_2, \\ \dot{z} =a\, z + 4\varepsilon\, H^3(z+g)\mathrm{Re}(\overline{\kappa}\,(z+g))\, 2x_2 - \varepsilon\, H^4 \frac{\partial g}{\partial X_2}\,\mathrm{Re}(\overline{\kappa}\,(z+g)). \end{cases} \tag{2.38}$$

Using the integral formula (2.3) for the function g we deduce that it is bounded, $g \leq C$. Since the function g prolongs to the (multivalued), holomorphic function ramified along the singular curve γ_0, the following bounds for derivatives of g hold

$$|g^{(k)}| \leq C_k |H|^{-k}. \tag{2.39}$$

Using this one can check that the system (2.38) satisfies the assumptions of Lemma 2.11. Thus, the invariant surface has the form

$$z = \frac{\varepsilon\, G - \varepsilon\, H^4 g}{\varepsilon\, H^4} = \varepsilon\, U(x,\varepsilon)$$

and the function U prolongs by zero to a C^1 function on a neighborhood of D. Thus, the function U satisfies the estimates

$$|U| \leq C|H|, \qquad |U'| \leq C,$$

which are equivalent to (2.36) and (2.37).

□

Now we show that the generalized Abelian integral $J(h)$ is a good approximation of the Poincaré return map and so the zeroes of $J(h)$ generate limit cycles for sufficiently small ε.

Proposition 2.15. *Let* $\Delta H(h,\varepsilon)$ *be the increment of the Hamiltonian after the first return of the system* V_ε *restricted to the invariant surface* L_ε*. Then, there exists a constant* C *such that*

$$|\Delta H - \varepsilon J| \leq C\,\varepsilon\,|h|^5, \tag{2.40}$$

$$|\partial_h(\Delta H) - \varepsilon J'(h)| \leq C\,\varepsilon\,|h|^4\ |\log h|. \tag{2.41}$$

Proof. Here we study the phase curves of the 2-dimensional vector field (2.31) i.e.

$$W_\varepsilon :\ = V_\varepsilon|_{L_\varepsilon} = X_H + \varepsilon Re(\overline{\kappa}\, G)\,\partial_{x_2}.$$

We fix the segment $I = \{(x_1, 0) : x_1 \in [-2,-1)\}$ transversal to the Hamiltonian flow. We denote by $\beta_\varepsilon(t,h)$ the integral curves of W_ε which start and finish at I. They satisfy

$$\begin{aligned} \dot{\beta}_\varepsilon(t,h) &= X_H + \varepsilon\,\mathrm{Re}(\overline{\kappa}\, G)\ e_2, \\ \beta_\varepsilon(0,h) &\in I, \qquad \beta_\varepsilon(T_\varepsilon,h) \in I \\ H(\beta_\varepsilon(0,h)) &= h. \end{aligned} \tag{2.42}$$

For $\varepsilon = 0$ the curve β_0 is the oval $\{H = h\}$ and for ε non-zero but small it is a small perturbation of β_0:

$$\beta_\varepsilon(t,h) = \beta_0(t,h) + \varepsilon b(t,h;\varepsilon).$$

Above $T_\varepsilon = T_\varepsilon(h)$ is the time of the first return to the unit I.

Lemma 2.16. *There exist a constant C and the positive, small constant ν such that the following estimates hold:*

$$|b| \le C|h|^4\, e^{(2\sqrt{3}+\nu)t}, \tag{2.43}$$

$$|\partial_h b| \le C|h|^3\, e^{(2\sqrt{3}+\nu)t}, \tag{2.44}$$

$$|T_\varepsilon - T_0| \le C\,\varepsilon\,|h|^3. \tag{2.45}$$

Proof. We use the scalar product $x \cdot x' = 3x_1\, x_1' + x_2\, x_2'$ and $|x| = \sqrt{x \cdot x}$.

It follows from the equation (2.42) that the function b satisfies the following initial value problem

$$\begin{cases} \dot{b} = \widetilde{\mathrm{d}X}_H\, b + \mathrm{Re}(\overline{\kappa}\, G(\beta_0 + \varepsilon\, b))\, e_2, \\ b(0,h;\varepsilon) = 0, \end{cases} \tag{2.46}$$

where $\widetilde{\mathrm{d}X}_H\, b = \frac{1}{\varepsilon}((X_H(\beta_0 + \varepsilon\, b) - X_H(\beta_0))$. We have $\widetilde{\mathrm{d}X}_H = \mathrm{d}X_H(\beta_0 + \theta\,\varepsilon\, b)$, for some $\theta \in (0,1)$. Hence

$$\widetilde{\mathrm{d}X}_H = \left(\begin{smallmatrix} 0 & -2 \\ -6x_1 & 0 \end{smallmatrix}\right), \qquad x_1 \in [-2,1].$$

Moreover, using estimates (2.36), (2.37), (2.39) we get $|G(\beta_0 + \varepsilon\, b)| \le C_1\, h^4$. For the solution b to the equation (2.46) we have

$$\begin{aligned} \tfrac{\mathrm{d}}{\mathrm{d}t}|b|^2 = 2|b|\, \tfrac{\mathrm{d}}{\mathrm{d}t}|b| &= 2b \cdot \widetilde{\mathrm{d}X}_H\, b + 2b \cdot \mathrm{Re}(\overline{\kappa}\, G(\beta_0 + \varepsilon\, b))\, e_2 = \\ &= -12(x_1+1)\, b_1\, b_2 + 2b_2\, \mathrm{Re}(\overline{\kappa}\, G(\beta_0 + \varepsilon\, b)) \le 4\sqrt{3}|b|^2 + 2C_2 h^4 |b|. \end{aligned}$$

Therefore $\frac{\mathrm{d}}{\mathrm{d}t}\,(|b|) \le 2\sqrt{3}|b| + C_2 h^4, \quad |b|(0) = 0$ and the Gronwall inequality [H] gives the bound (2.43).

Since the difference of flows $\varepsilon\, b$ after the Hamiltonian period T_0 is $\le \widetilde{C}|h|^3$ and the "velocity" $|V_\varepsilon| \sim 1$, the difference of periods $|T_\varepsilon - T_0|$ satisfies (2.45).

The derivative $\frac{\partial b}{\partial_h}$ satisfies the respective linear variation equation related to (2.46)

$$\tfrac{\mathrm{d}}{\mathrm{d}t}\left(\tfrac{\partial b}{\partial_h}\right) = \left(\widetilde{\mathrm{d}X}_H + \varepsilon\,(\ldots)\right) \tfrac{\partial b}{\partial_h} + h^3\,(\ldots)\,(\tfrac{\partial \beta_0}{\partial_h}), \qquad \tfrac{\partial b}{\partial_h}(0,h;\varepsilon) = 0,$$

where we denoted by $(\ldots)$ the bounded terms (see estimations (2.43,2.45,2.37,2.39)). Since the flow variation $\frac{\partial \beta_0}{\partial_h}$ of the Hamiltonian field satisfies $\left|\frac{\partial \beta_0}{\partial_h}\right| \le C\, e^{2\sqrt{3}s}$, the inequality (2.44) holds. This finishes the proof of Lemma 2.16.

□

We continue the proof of Proposition 2.15.

We split the difference between the Poincaré map and the linearization $\varepsilon\, J(h)$ in two integrals $R_1(h,\varepsilon)$ and $R_2(h,\varepsilon)$:

$$\begin{aligned} R_1 &= \int_{\gamma_\varepsilon} \mathrm{Re}(\overline{\kappa}\,(G - H^4 g))\, \mathrm{d}x_1, \\ R_2 &= \int_{\gamma_\varepsilon} \mathrm{Re}(\overline{\kappa}\, H^4 g)\, \mathrm{d}x_1 - \int_{\gamma_0} \mathrm{Re}(\overline{\kappa}\, H^4 g)\, \mathrm{d}x_1. \end{aligned}$$

We shall show that the estimations (2.40) and (2.41) hold for both R_1 and R_2.

The inequality (2.40) is a direct consequence of the bound (2.36). The difference of R_1 in close values of h takes the form

$$R_1(h+\delta)-R_1(h)=\int_{\gamma_\varepsilon(h,h+\delta)}\partial_{x_2}\mathrm{Re}(\overline{\kappa}\,(G-H^4g))\,\mathrm{d}x_2\wedge\mathrm{d}x_1+O(\varepsilon|h-2|^5),$$

where the integral is taken along the strip $\gamma_\varepsilon(h,h+\delta)$ between $\gamma_\varepsilon(h)$ and $\gamma_\varepsilon(h+\delta)$. The area of this strip is of the same order as the area of the domain $\{X:\ h<H<h+\delta\}$, i.e. $\sim\delta\cdot I_0\sim C\,\delta\,|\log h|$ (see proof of Lemma 2.2). So the estimate (2.41) follows (2.37).

To prove the estimate for R_2 we use (2.42) and (2.39):

$$|R_2|\le C_1\int_0^{T_0}\mathrm{Re}(\overline{\kappa}\,H^4g)(\beta_0+\varepsilon b)-\mathrm{Re}(\overline{\kappa}\,H^4g)(\beta_0)+\int_{T_0}^{T_\varepsilon}C_2\,\varepsilon\,|h|^4\le$$
$$\le C_3\,\varepsilon\,|h|^7\int_0^{T_0}e^{(2\sqrt{3}+\nu)t}\,\mathrm{d}t+C_4\,|h|^4\,|T_\varepsilon-T_0|\le C_5\,\varepsilon\,|h|^{6-2\nu}\le C\varepsilon|h|^5.$$

Similarly, using the following formula for differential of integral

$$\frac{\partial}{\partial h}\int_{\beta_\varepsilon}\omega=\int_{\beta_\varepsilon}i_{\partial\beta_\varepsilon/\partial_h}\,\mathrm{d}\omega+\omega\left(\left.\frac{\partial\beta_\varepsilon}{\partial_h}\right|_{t=0}\right)-\omega\left(\left.\frac{\partial\beta_\varepsilon}{\partial_h}\right|_{t=T_\varepsilon}\right)$$

and the bounds (2.42), (2.39), we get

$$|R_2|\le C\,\varepsilon\,|h|^5,\qquad|\partial_hR_2|\le C\,\varepsilon\,|h|^4.$$

□

Now we can finish the proof of the Main Theorem. The restriction of system (1.3) to its invariant surface $L_\varepsilon=\{y=\varepsilon\,G(x,\varepsilon)\}$ has the form (2.31). The increment $\Delta H=\mathcal{P}(h)-h$, associated with the Poincaré map $\mathcal{P}(h)$ (on a section transversal to γ_0), equals (see Proposition 2.15)

$$\Delta H(h)=\varepsilon J(h)+O(|\varepsilon|\,|h|^5),$$
$$(\Delta H)'(h)=\varepsilon J'(h)+o(|\varepsilon|\,|h|^{4-1/4}).$$

Since we also have (see proof of Corollary 2.5)

$$J(h)=Rh^{4+1/2}\cos(\log\sqrt{h}-\alpha_0)+o(h^{4-1/4}),$$

any simple zero of $J(h)$, which is sufficiently close to 0 generates, by the Implicit Function Theorem, a simple zero of ΔH. Thus the sequence $h_n\to0^+$ of simple zeroes of $J(h)$ (see Corollary 2.5) guarantees the existence of *infinite sequence* $\widetilde{h}_n\to0^+$, $n\ge N_0$ of simple zeroes of the increment ΔH. Any simple zero of ΔH corresponds to a limit cycle of (1.3).

The proof of Main Theorem is now complete.

□

References

[AI] Arnold V. I. and Il'yashenko Yu. S., *Ordinary differential equations, in: "Ordinary Differential Equations and Smooth Dynamical Systems"*, Springer–Verlag, New York, (1997), pp. 1–148; (Russian: Fundamental Directions, v. **1**, VINITI, Moscow, 1985, pp. 1–146).

[Be] Belitskii H., "Normal forms, invariant and local mappings", Naukova Dumka, Kiev, 1979 (in Russian).

[Bo] Bobieński M., *Contour integrals as* Ad*-invariant functions on the fundamental group* (submitted).

[BZ1] Bobieński M. and Żołądek H., *Limit cycles for multidimensional vector field. The elliptic case*, J. Dynam. Control Systems, **9**, (2003), No 2, 265–310.

[BZ2] Bobieński M. and Żołądek H., *Limit cycles of three dimensional polynomial vector fields*, Nonlinearity **18**, (2005), No 1, 175–209.

[E] Ecalle J., "Introduction aux fonctions analysables et preuve constructivede la conjecture de Dulac", Actualités Mathématiques, Herman, Paris, 1992.

[GR] Gradshteyn I. S., Ryzhik I. M., "Table of Integrals, Series and Products. Fifth edition", Academic Press, Inc. 1994.

[GuHo] Guckenheimer J. and Holmes P., "Nonlinear oscillations, dynamical systems, and bifurcations of vector fields", Applied Mathematical Sciences, **42**, Springer-Verlag, 1983.

[H] Hartman, P., "Ordinary differential equations", Classics in Applied Mathematics, **38**, Philadelphia, PA, 2002.

[HPS] Hirsch M., Pugh C., Shub M., "Invariant manifolds", Lect. Notes in Math. **583**, Springer-Verlag, New York, 1977.

[Il] Il'yashenko Yu. S., *Centennial history of Hilbert's 16th problem*, Bull. Amer. Math. Soc. **39** (2002), No 3, 301–354.

[M] Melnikov V. K., *On the stability of a center for time-periodic perturbations*, Trans. Moscow Math. Soc. **12** (1963), 1–57.

[MiMr] Mischaikow K., Mrozek M., *Chaos in the Lorenz equations: a computer-assisted proof*, Bull. Amer. Math. Soc. (N.S.), **32**, (1995), No 1, 66–72.

[Ni] Nitecki Z., "Differentiable dynamics. An introduction to the orbit structure of diffeomorphisms", MIT Press, Cambridge, 1971.

[P] Pontryagin L. S., *On dynamical systems close to Hamiltonian systems*, in: "Selected Works", v. 1, Gordon & Breach, New York, 1986; [Russian: Zh. Ekper. Teoret. Fiziki 4 (1934), 234–238].

[LZ] Leszczyński P., Żołądek H., *Limit cycles appearing after perturbation of certain multidimensional vector fields*, J. Dynam. Diff. Equat. **13** (2001), No 4, 689–709.

[Yak] S. Yakovenko, *On functions and curves defined by ordinary differential equations*, The Arnoldfest (Toronto, ON, 1997), Fields Inst. Commun., **24**, Amer. Math. Soc., Providence, 1999, pp. 497–525.

[Z2] H. Żołądek, "The Monodromy Group", Monografie Matematyczne, Birkhäuser, Basel, 2006.

INSTITUTE OF MATHEMATICS, WARSAW UNIVERSITY, UL. BANACHA 2, 02-097 WARSAW, POLAND

E-mail address: `mbobi@mimuw.edu.pl`

E-mail address: `zoladek@mimuw.edu.pl`